

Client Information

<i>Client Name</i>	Client Demo
<i>Date of birth (age)</i>	18 January 1998 (22)

Assessment Information

<i>Assessment</i>	Patient Health Questionnaire - Depression (PHQ-9)
<i>Date administered</i>	10 October 2020
<i>Assessor</i>	Mr Demo Assessor
<i>Time taken</i>	1 minutes 26 seconds

Results

	Score	Value
		22
	Percentile (non-depressed)	99.9
	Percentile (major depression)	78.9
	Percentile (other depression)	98.4

Interpretive Text

<p>This individual scored as having a severe level of depression. This individual scored above 10, indicating that this individual may have Major Depression, with a sensitivity of at least 88% and a specificity of 88% This person reported that they thought about death or hurting themselves several days over the last two weeks.</p>

Scoring and Interpretation Information

<p>A raw score (from 0 to 27) is presented, as well as a percentile based on sample of non-depressed individuals, a sample of those diagnosed with major depression, and sample diagnosed as having other depressive disorders.</p> <p>An individual with a percentile of around 50 compared to the major depression (MD) sample has a typical level of depression when compared to others diagnosed with major depression. The non-depressed percentile and the other-depressed percentile will be substantially higher. Conversely, an individual who scores in the 50th percentile compared to the non-depressed sample will likely score very low on the two depressed sample subscales.</p>
--

Client Responses

		Not at all	Several days	More than half the days	Nearly every day
1	Little interest or pleasure in doing things	0	1	2	3

Client Name	Client Demo
--------------------	-------------

Client Responses (cont.)		Not at all	Several days	More than half the days	Nearly every day
2	Feeling down, depressed, or hopeless	0	1	2	3
3	Trouble falling or staying asleep, or sleeping too much	0	1	2	3
4	Feeling tired or having little energy	0	1	2	3
5	Poor appetite or overeating	0	1	2	3
6	Feeling bad about yourself — or that you are a failure or have let yourself or your family down	0	1	2	3
7	Trouble concentrating on things, such as reading the newspaper or watching television	0	1	2	3
8	Moving or speaking so slowly that other people could have noticed? Or the opposite — being so fidgety or restless that you have been moving around a lot more than usual	0	1	2	3
9	Thoughts that you would be better off dead or of hurting yourself in some way	0	1	2	3