

Client Information

<i>Client Name</i>	Demo Client
<i>Date of birth (age)</i>	14 February 1961 (59)

Assessment Information

<i>Assessment</i>	Opioid Risk Tool (ORT)
<i>Date administered</i>	12 January 2019
<i>Assessor</i>	Mr Demo Assessor
<i>Time taken</i>	0 minutes 8 seconds

Results

	Value
Total Opioid Risk score	12

Interpretive Text

<p>This individual scores in the high risk category (scores more than 8), indicating that their risk of engaging in aberrant behaviours is approximately 91 per cent.</p>

Scoring and Interpretation Information

<p>Scores range from 0 to 26, with scores calculated differently dependent on gender and age. The scores are categorized into three risk categories.</p> <ul style="list-style-type: none"> - Low Risk= Scores 0-3 - Moderate Risk = scores 4 – 7 - High risk = Scores 8 or more <p>According to the validation sample (Webster, 2005) patients in the low risk category have a 5.6% chance of developing one or more aberrant behaviour when prescribed opioids. The moderate risk category had a risk of 28% while the high risk category had a risk of 91%. Note that these percentages are not direct measures of risk of opioid abuse or dependence, but rather aberrant behaviours (which are known to be associated with abuse and dependence). Aberrant behaviours are defined as:</p> <ul style="list-style-type: none"> -Used additional opioids than those prescribed -Used additional opioids than those prescribed more than once -Forged prescription -Sold prescription -Admitted to seeking euphoria from opioids -Admitted to wanting opioids for anxiety -Overdose and death -Injected drug -Abnormal urine/blood screen -Abnormal urine/blood screen positive for 2 or more

Client Name	Demo Client
--------------------	-------------

Scoring and Interpretation Information (cont.)

- substances
- Solicited opioids from other providers
- Unauthorized ER visits
- Concurrent abuse of alcohol
- Unauthorized dose escalation

Client Responses

		Yes	No
1	Has there been family history of alcohol abuse?	Yes	No
2	Has there been family history of illegal drug use?	Yes	No
3	Has there been family history of recreational drug use?	Yes	No
4	Has there been personal history of alcohol abuse?	Yes	No
5	Has there been personal history of illegal drug use?	Yes	No
6	Has there been personal history of recreational drug use?	Yes	No
7	Aged between 16 - 45 years?	Yes	No
8	Has there been a history of preadolescent sexual abuse?	Yes	No
9	Has there been a personal history of Attention Deficit Disorder (ADD or ADHD), bipolar or schizophrenia?	Yes	No
10	Has there been a personal history of depression?	Yes	No