

### Client Information

<i>Client Name</i>	Dummy Client
<i>Date of birth (age)</i>	1 January 2000 (24)

### Assessment Information

<i>Assessment</i>	Obsessional Compulsive Inventory – Revised (OCI-R)
<i>Date administered</i>	10 February 2024
<i>Assessor</i>	Mr Jimmy Whispers
<i>Time taken</i>	0 minutes 18 seconds

### Results

	Score	Normative Percentile	Clinical Percentile
OCD	28	99.9	63.1
Hoarding Disorder	6	98.5	9

### Interpretive Text

This OCD score reached the cutoff (greater than or equal to 12), indicating the responses are consistent with OCD with a sensitivity of 82% and specificity of 83% (Wootton et al., 2015). A score of 12 or greater correctly identified 83% of those diagnosed with OCD in the clinical sample (Wootton et al., 2015).


This hoarding disorder score reached the cutoff (greater than or equal to 6), indicating the responses are consistent with hoarding disorder with a sensitivity of 92% and specificity of 93% (Wootton et al., 2015). A score of 6 or greater correctly identified 93% of those diagnosed with hoarding disorder in the clinical sample (Wootton et al., 2015).

### OCD Subscales


	Raw Score (0-12)	Clinical Percentile
Washing	4	44.3
Obsessing	5	30.8
Ordering	9	91.3
Checking	5	60.1
Neutralising	5	73.9

<b>Client Name</b>	Dummy Client
--------------------	--------------

**OCD Score**



**Hoarding Score**



<b>Scoring and Interpretation Information</b>
For the OCD component of the OCI-R (items 2, 3, 4, 5, 6, 8, 9, 10, 11, 12, 14, 15, 16, 17, 18), the total score ranges from 0 – 60, with

<b>Client Name</b>	Dummy Client
--------------------	--------------

**Scoring and Interpretation Information (cont.)**

higher scores indicative of more severe OCD symptoms. A cutoff score of 12 is used to determine the likelihood of an OCD diagnosis (with a sensitivity of 82% and specificity of 83%).

Normative and clinical percentiles are presented comparing the respondent's scores to other adults (Wootton et al., 2015). A normative percentile rank of 50 indicates an average level of OCD symptoms in comparison to the the general population, and is indicative of typical (and healthy) levels of symptomatology. A clinical percentile rank of 50 indicates an average level of OCD symptoms in comparison to the clinical group (with an OCD diagnosis), and is indicative of elevated levels of symptomatology.

For the hoarding disorder subscale of the OCI-R (items 1, 7, 13), the total score ranges from 0 – 12, with higher scores indicative of more severe hoarding symptoms. A cutoff score of 6 is used to determine the likelihood of a hoarding disorder diagnosis (with a sensitivity of 92% and specificity of 93%).

A normative and clinical percentile are presented comparing the respondent's scores to other adults (Wootton et al., 2015). A normative percentile rank of 50 indicates an average level of hoarding symptoms in comparison to the normative group, and is indicative of typical (and healthy) levels of symptomatology. A clinical percentile rank of 50 indicates an average level of OCD symptoms in comparison to the clinical group (with a hoarding disorder diagnosis), and is indicative of elevated levels of symptomatology.

The OCD component of the OCI-R also reports the client's score (between 0 – 12) across 6 subscales, with a clinical percentiles comparing the respondent's scores to a comparison group whom have received a OCD diagnosis (Abramovitch et al., 2020):

Washing (items 5, 11, 17) – assessing difficulty in touching objects that have been touched before and excessive washing due to feeling contaminated.

Obsessing (items 6, 12, 18) – assessing difficulty with thoughts including trying to control them, becoming upset by unpleasant thoughts, and a feeling of excessive unpleasant thoughts.

Ordering (items 3, 9, 15) – assessing challenges with ordering of objects.

Checking (items 2, 8, 14) – assessing excessive checking of items (doors, windows, drawers, taps, switches).

Neutralising (items 4, 10, 16) – assessing compulsions to count and excessive feelings towards numbers.

<b>Client Name</b>	Dummy Client
--------------------	--------------

<b>Client Responses</b>		Not at all	A little	Moderately	A lot	Extremely
1	I have saved up so many things that they get in the way.	0	1	2	3	4
2	I check things more often than necessary.	0	1	2	3	4
3	I get upset if objects are not arranged properly.	0	1	2	3	4
4	I feel compelled to count while I am doing things.	0	1	2	3	4
5	I find it difficult to touch an object when I know it has been touched by strangers or certain people.	0	1	2	3	4
6	I find it difficult to control my own thoughts.	0	1	2	3	4
7	I collect things I don't need.	0	1	2	3	4
8	I repeatedly check doors, windows, drawers, etc.	0	1	2	3	4
9	I get upset if others change the way I have arranged things.	0	1	2	3	4
10	I feel I have to repeat certain numbers.	0	1	2	3	4
11	I sometimes have to wash or clean myself simply because I feel contaminated.	0	1	2	3	4
12	I am upset by unpleasant thoughts that come into my mind against my will.	0	1	2	3	4
13	I avoid throwing things away because I am afraid I might need them later.	0	1	2	3	4
14	I repeatedly check gas and water taps and light switches after turning them off.	0	1	2	3	4
15	I need things to be arranged in a particular way.	0	1	2	3	4
16	I feel that there are good and bad numbers.	0	1	2	3	4
17	I wash my hands more often and longer than necessary.	0	1	2	3	4
18	I frequently get nasty thoughts and have difficulty in getting rid of them.	0	1	2	3	4